Shurguard Plus

U S Chemica

Read the label affixed to the container for this product before applying. Use of this product according to the reference sheet is subject to the use precautions and limitations imposed by the label affixed to the container for Shurguard Plus. For use in hospitals, nursing homes, hotels, food service establishments, restaurants and bars where disinfection, sanitization and deodorization is of prime importance. When used as directed, this product is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms cabinets, tables and chairs. Effective against the following pathogens on hard, non-porous surfaces:

BACTERIA	Listeria monocytogenes Pseudomonas aeruginosa Salmonella entericia Staphylococcus aureus	Vancomycin intermediate resistant Staphylococcus aureus (VISA) Yersinia enterocolitica	FOOD CONTACT SANITIZATION	Campylobacter jejuni Escherichia coli Escherichia coli 0157:H7 Klebsiella penumoniae Listeria monocytogenes Methicillin resistant Staphylococcus aureus (MRSA)	Salmonella typhi Shigella sonnei Staphylococcus aureus Vancomycin resistant Enterococcus faecalis (VRE) Vibrio cholera Yersinia enterococlitica
ហ	*Adenovirus Type 5 *Hepatitis B virus (HBV) *Hepatitis C virus (HCV)	*Laryngotracheitis virus * SARS-Related Coronavirus 2 (SARS-CoV-2)1			
VIRUSES	*Herpes Simplex virus Type 1 *Human coronavirus *HIV-1 (AIDS virus) *Human Influenza virus *Influenza A virus (H7N9) *Influenza A2	*SARS Associated Coronavirus *Vaccinia virus ¹ SARS-CoV – 2 fl. oz. per gal. of water for 30 seconds.	ANIMAL	*Avian Influenza A *Bovine Viral Diarrhea virus (BVDV) *Newcastle Disease virus	*Procine Respiratory & Reproductive Syndrome virus (PRRSV)
DILUTION	General Disinfection – 3 fl. oz. per 5 gallons of water Hospital Disinfection – 3.5 fl. oz. per 5 gallons of water Virucidal* Activity – 3.5 fl. oz. per 5 gallons of water Mildewstat – 3 fl. oz. per 5 gallons of water		NFC SANITIZER	Klebsiella pneumoniae Listeria monocytogenes Staphylococcus aureus	
סורח	Non-Food Contact (NFC) Sanitization – 0.25 fl. oz. per 1 gallon of water Food Contact Sanitization – 0.25 - 0.5 fl. oz. per 1 gal of water		CTIVE EDIENTS	Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium5.0% Alkyl (68% C ₁₂ , 32% C ₁₂) dimethyl ethylbenzyl ammonium	
	KEEP OUT OF REACH OF CHILDREN DANGER			Chloride	

ADDITIONAL DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Labeling must be in the possession of the user at the time of using the product. See container label for FIRST AID, storage and Disposal instructions and PRECAUTIONARY STATEMENTS and complete DIRECTIONS for USE before applying. Use of this product is subject to the precautions and limitations imposed by the label affixed to the container for this product.

USE SITES: Beauty Salons, Ships, Vehicles, Diary Farms, Animal Kennels, Dairy Processing Plants, Automotive Garages, Meat Processing Establishments, Education Institutions, Animal Care Facilities, Patient Care Facilities, Physician Offices, Police Stations, Performance Centers, Airplanes, Animal Life Science Laboratories, Auto Repair Centers, Classrooms, EMS & Fire Facilities, Factories, Food Processing Plants, Health Clubs, Institutional Facilities, Laboratories, Life Care Retirement Communities, Lodging Establishments, Medical Offices, Medical Research Facilities, Mortuaries, Municipal Government Buildings, Nurseries, Nursing Homes, Post Offices, Recreational Facilities, Recycling Centers and Equipment, Schools, Shopping Malls, Spas, Supermarkets, Transporation Terminals, Veterinary Clinics, Zoos. For a complete list of use sites and uses, consult your sales representative.

NON-FOOD CONTACT SURFACE SANITIZATION DIRECTIONS:

Sanitization will occur only on pre-cleaned surfaces. The sanitizer use solution can be applied to pre-cleaned, hard, inanimate, non-porous surfaces with a cloth, sponge, mop, sprayer or by flooding or immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from the surface. Rub with brush, sponge or cloth. Do not breathe spray.

Treated surfaces must remain visibly wet to 30 seconds and allowed to air dry.

NOTE: With spray applications, cover or remove all food products.

Non-Acid Toilet Bowl and Urinal Disinfectant Directions: Remove visible soil prior to disinfecting. Empty toilet bowl or urinal and apply a use-solution of 3.5 fl. oz. per 5 gal of water to exposed surfaces including under the rim with a cloth, mop, sponge or hand pump trigger sprayer. Brush or swab thoroughly, then allow to remain visibly wet for 10 minutes and flush.

To Disinfect Food Processing Premises: Floors, walls, and storage areas, add 3 fl. oz. of this product per 5 gallons of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes, then remove excess liquid. Before using this product, food products and packaging materials must be removed from the area or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

Drain Cleaning: For cleaning drains, dilute 1.25 oz per 1 gal of water. Pour solution down drain being sure to coat all sides of drainpipe. Allow all treated surfaces exposed to solution to air dry.

Sanitization of Beverage Dispensing Equipment: Wash equipment with a compatible detergent and rinse with potable water prior to sanitizing. Fill equipment with a use solution of 0.25 - 0.5 fl. oz. of this product per gal of water. Allow use solution to remain in equipment for at least 60 seconds. Drain thoroughly and allow to air dry before reuse. Do not rinse with potable water.

SKU 4/1 gal. (057602), 5 gal. pail (3296521) and 55 gal. drum (073157)